

DIRECT MEASUREMENT OF THE RADIUS AND DENSITY OF THE TRANSITING EXOPLANET HD 189733b WITH THE CHARA ARRAY

ELLYN K. BAINES,¹ GERARD T. VAN BELLE,² THEO A. TEN BRUMMELAAR,¹ HAROLD A. MCALISTER,¹ MARK SWAIN,³
NILS H. TURNER,¹ LASZLO STURMANN,¹ AND JUDIT STURMANN¹

Received 2007 February 23; accepted 2007 April 20; published 2007 May 9

ABSTRACT

We have measured the angular diameter of the transiting extrasolar planet host star HD 189733 using the CHARA optical/IR interferometric array. Combining our new angular diameter of 0.377 ± 0.024 mas with the *Hipparcos* parallax leads to a linear radius for the host star of $0.779 \pm 0.052 R_{\odot}$ and a radius for the planet of $1.19 \pm 0.08 R_{\text{Jup}}$. Adopting the mass of the planet as derived by its discoverers, we derive a mean density of the planet of $0.91 \pm 0.18 \text{ g cm}^{-3}$. This is the first determination of the diameter of an extrasolar planet through purely direct means.

Subject headings: infrared: stars — stars: fundamental parameters — stars: individual (HD 189733) — techniques: interferometric

1. INTRODUCTION

A handful of extrasolar planets transit their host stars, causing a reduction in stellar flux as the planet blocks part of the star's disk. The planet orbiting HD 189733 is one of the 14 known transiting planets. Using radial velocity and photometric measurements made at the Haute-Provence Observatory, Bouchy et al. (2005) discovered a hot Jupiter-like planet with an orbital period of 2.219 days and estimated the star's radius to be $0.76 \pm 0.01 R_{\odot}$. This value, along with a planet-to-star radius ratio of 0.172 ± 0.003 , led to a planetary radius of $1.26 \pm 0.03 R_{\text{Jup}}$. More recently, Bakos et al. (2006b) refined the orbital parameters using *BVRI* multiband photometry and found the planet's radius to be $1.154 \pm 0.032 R_{\text{Jup}}$.

We observed HD 189733 using Georgia State University's Center for High Angular Resolution Astronomy (CHARA) Array in order to directly determine the host star's radius and thereby calculate, in a strictly geometric manner, the radius and density for the planet.

Planetary densities were previously estimated from photometric observations of the transiting planets and range from 0.38 g cm^{-3} for HD 209458b (Charbonneau et al. 2000) to 1.17 g cm^{-3} for HD 149026b (Sato et al. 2005). These density calculations are highly dependent on estimated stellar diameters based on spectral energy distribution (SED) fits using published photometric values, which are fundamentally indirect in nature, relying upon a priori assumptions regarding the host stars' stellar atmospheres. For the four "bright" ($V < 12$) transit host stars, these angular sizes are in the range of 0.05–0.40 mas. The longest baselines of the CHARA Array are capable of resolving the largest and brightest of these objects.

2. INTERFEROMETRIC OBSERVATIONS AND DIAMETER DETERMINATION

2.1. Observations and Data Reduction

Spatially resolved observations of HD 189733 were obtained with the CHARA Array, a six-element interferometer located on

Mount Wilson, California (McAlister et al. 2005). The Array operates in two wavelength regimes: in visible wavelengths (470–800 nm) for tracking and tip/tilt correction, and in the near-infrared *K'* (2.13 μm) and *H* (1.67 μm) bands for fringe detection. Because of the small angular diameter for the star, only the *H*-band observations obtained at our longest baseline pair (telescopes E1 and S1) are used in our final diameter analysis.

HD 189733 was observed on several nights during the summer of 2006 along with the calibrator star HD 190993, a B3 V star offset by 1.7° , selected on the basis of its small estimated angular diameter and its apparent lack of any close companion. The latter criterion was verified by a thorough literature search, while a SED fit to HD 190993 led to an estimated angular diameter of 0.167 ± 0.035 mas with no residuals suggestive of a companion (see Fig. 1). This results in a predicted visibility (V) for the calibrator of $V_{\text{cal}} = 0.961^{+0.019}_{-0.008}$ at our longest baseline of 330 m, resulting in a contribution of $\sigma_v \approx 0.01$ –0.02 to the calibrated visibility errors seen in Table 1. The small angular size and high visibility of the calibrator mean that HD 190993 is essentially unresolved using the CHARA Array, and the uncertainty in visibility due to calibrator diameter error is small compared to the measurement error. Therefore, uncertainties in the calibrator diameter will not affect the HD 189733 diameter measurement significantly (van Belle & van Belle 2005). Even HD 190993's considerable $v \sin i$ does not contribute an error to our diameter fits due to its small angular size.

We note that the M dwarf companion to HD 189733 reported by Bakos et al. (2006a) on the basis of common space motion at an angular separation of 11.2" is well outside the interferometric field of view, and its presence has no effect on our results. Although the effect on visibility would be small in the first lobe of the $V(B)$ curve, we have confirmed that our observed epochs do not occur within the predicted times of planetary transit or eclipse using the period and reference time of central transit of Bakos et al. (2006b).

All our observations were obtained with the single-baseline, pupil-plane "CHARA Classic" beam combiner, and we employed the standard practice of observing the target and calibrator sequentially to provide a series of time-bracketed observations from which the instrumental visibilities could be reduced to calibrated values for the target star. The observing practice and reduction process employed here are identical to

¹ Center for High Angular Resolution Astronomy, Georgia State University, P.O. Box 3969, Atlanta, GA 30302-3969; baines@chara.gsu.edu, theo@chara-array.org, hal@chara.gsu.edu, niels@chara-array.org, sturmann@chara-array.org, judit@chara-array.org.

² Michelson Science Center, California Institute of Technology, 770 South Wilson Avenue, MS 100-22, Pasadena, CA 91125; gerard@ipac.caltech.edu.

³ Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena, CA 91109; mark.swain@jpl.nasa.gov.




FIG. 1.—SED fit of our calibrator star, HD 190993. In the top panel, the vertical bars represent the errors for the data points that they overlay, and the horizontal bars represent the bandpass of the data points. In the bottom panel, the fractional residuals (difference between data point and fit point, normalized by that data point) are shown for each data point.

those described by ten Brummelaar et al. (2005). The results of this process are summarized in Table 1.

Single-baseline Michelson stellar interferometers measure complex visibilities, usually recorded as amplitudes and phases, which are related to the intensity distribution of the target through a Fourier transform; phase information is typically corrupted by the atmosphere, leaving the amplitude, referred to simply as the *visibility* (Colavita et al. 1999).

2.2. Diameter Fit

Diameter fits to the visibilities and baselines from Table 1 were performed using the uniform-disk (UD) approximation given by

$$V = \frac{2J_1(x)}{x}, \quad (1)$$

where J_1 is the first-order Bessel function and

$$x = \pi B \theta_{\text{UD}} \lambda^{-1}, \quad (2)$$

where B is the projected baseline at the star's position, θ_{UD} is the apparent UD angular diameter of the star, and λ is the wavelength of the observation. The limb-darkened (LD) relationship incorporating the linear limb-darkening coefficient μ_λ (Hanbury Brown et al. 1974) is given by

$$V = \left(\frac{1 - \mu_\lambda}{2} + \frac{\mu_\lambda}{3} \right)^{-1} \left[(1 - \mu_\lambda) \frac{J_1(x)}{x} + \mu_\lambda \left(\frac{\pi}{2} \right)^{1/2} \frac{J_{3/2}(x)}{x^{3/2}} \right]. \quad (3)$$

These fits resulted in $\Theta_{\text{UD}} = 0.366 \pm 0.024 \text{ mas}$ and $\Theta_{\text{LD}} = 0.377 \pm 0.024 \text{ mas}$, the latter incorporating $\mu_\lambda = 0.36$ taken from Claret et al. (1995) after adopting $\log g = 4.5$ and $T_{\text{eff}} = 5000 \text{ K}$ for HD 189733 (see Fig. 2). The reduced χ^2 minimization in both cases yielded a value of 1.593, and the errors quoted are for an increase of the χ^2 value of 1.0, that is, the 68% confidence interval. Dividing this χ^2 by the number of degrees of freedom, which in our case is 8, yields 0.199, which is much less than 1.0, showing that the fit is quite good

TABLE 1
INTERFEROMETRIC MEASUREMENTS OF HD 189733

MJD (53,886.0+)	Baseline (m)	Visibility	σ_V
0.905	330.5	0.851	0.071
1.936	327.9	0.843	0.056
1.958	324.9	0.857	0.054
8.865	330.5	0.869	0.034
76.742	326.5	0.909	0.069
76.761	323.8	0.863	0.049
76.778	321.3	0.877	0.045
76.793	319.0	0.839	0.045
76.824	315.5	0.829	0.061

and that our error estimates for the visibility points are conservative. If we rescale these error bars to force χ^2 to be equal to the number of degrees of freedom, which assumes that there are no systematics in the measurements, they are approximately half the size that they are shown in Figure 2 and would also reduce our final error estimates by a factor of 2. However, we will remain conservative and continue to use the error estimate based on the raw χ^2 value.

2.3. Estimate of the Angular Size of HD 189733

An a priori estimate of the angular size of HD 189733 is a parameter of considerable interest, because the size of HD 189733b is determined only relative to the size of its parent star from the photometric transit timing data. Bakos et al. (2006b) consider no less than four separate methods in their investigation of the system: the $V - K$ color angular radius prediction (Kervella et al. 2004), the temperature radius, isochrone radii from Girardi et al. (2002) and Baraffe et al. (1998), and the Johnson $V-2\text{MASS } T_{\text{eff}}$ calibration of Masana et al. (2006).

None of these approaches appear to have much merit, since the only primary data we have been able to find in the literature were Tycho B_T , V_T (Bessell 2000), Strömgren $ubvy$ (Olsen 1993), and 2MASS JHK photometry (Cutri et al. 2003). Neither spectroscopy nor measures of $\log g$ nor direct measures of Johnson photometry appear to be available in the literature. The values of V used in Kervella et al. (2004) and Masana et al. (2006) appear to have been extrapolated from V_T . Furthermore, sustained long-term observations of HD 189733 by the *MOST* asteroseismology satellite have found the star to be pho-




FIG. 2.—The χ^2 fit to HD 189733's visibilities. The solid line represents a theoretical visibility curve for a star with a limb-darkened diameter of 0.377 mas, the squares are the measured visibilities, and the vertical lines are the measured errors.

ometrically variable (J. Rowe 2007, private communication), casting significant doubt on any radius derived from a photometric relationship. Based on this information, we consider the size errors for HD 189733 quoted in Bouchy et al. (2005) and Bakos et al. (2006b), derived from the methods cited above, to be underestimates.

To explore what is a more appropriate error for an inferred angular size, we executed a SED fit of the available spectrophotometry for HD 189733 cited above. Given the known variability of HD 189733, the quoted millimagnitude error estimates of the Tycho and Strömgren photometric data points were increased by a factor of 10. These photometric data points were fit to the solar-abundance K0 V and K2 V templates available from Pickles (1998), with the resulting fit values for reddening, bolometric flux, and angular diameter seen in Table 2, along with the appropriate χ^2 per degree of freedom (χ^2 PDF) values. Unfortunately, a K1 V template is not available in that library, although an estimate of one from interpolating between the two bracketing spectral types was synthesized by us for testing this spectral type. These fits are seen in Figure 3.

The appropriate model spectrum from Munari et al. (2005) for a 5000 K star was fit with a χ^2 PDF of 2.80, but this model (and the others available in the literature) unfortunately only covered the visible portion of the spectrum. The 115–2500 nm range of Pickles (1998) was necessary to fully characterize the available photometry, and thus we constrained our analysis to this particular set of stellar templates.

Our finding is that, even with this highly detailed analysis of the stellar SED, the most appropriate modeling of that SED reveals a predicted angular size of only $\theta = 0.363 \pm 0.011$ mas—a 3% error bar—which corresponds to a stellar linear radius of $R = 0.752 \pm 0.026 R_{\odot}$.

3. DISCUSSION

Our new direct determination for the angular diameter of HD 189733 of $\theta_{LD} = 0.377 \pm 0.024$ mas can be combined with the *Hipparcos* parallax for the star of $\pi = 51.9 \pm 0.9$ mas (Perryman et al. 1997) to give a physical radius for the star of $R_{\text{star}} = 0.779 \pm 0.052 R_{\odot}$, which is about 3% larger than that adopted by Bakos et al. (2006b).

By the nature of the light-curve analysis, the relative increase in the radius of the host star will directly translate into the same relative increase in the radius of the planet HD 189733b. Thus, revising the radius of Bakos et al. (2006b) of $1.154 R_{\text{Jup}}$, our new estimate for this value is $R_{\text{planet}} = 1.19 \pm 0.08 R_{\text{Jup}}$. Furthermore, adopting the value of Bouchy et al. (2005) for the mass of the planet of $1.15 M_{\text{Jup}}$, we derive a new estimate for the density of HD 189733b, $\rho = 0.91 \pm 0.18 \text{ g cm}^{-3}$. These values are in good agreement with Winn et al. (2007), who used transit photometry to constrain the stellar and planetary radii. The values of M_{planet} , R_{planet} , and ρ_{planet} are all consistent with the modest collection of these parameters presently available for transiting exoplanet systems, and they support the conclusion that HD 189733 is not among the few hot Jupiters that present extraordinarily large radii for their masses.

4. INTERFEROMETRIC NONDETECTION OF BINARITY OF HD 189733

Given the higher resolution of interferometric arrays, a possible close-separation tertiary companion may affect our measures of HD 189733's visibility and thereby complicate our interpretation. As such, it was prudent for us to also observe

HD 189733 with the Palomar Testbed Interferometer (PTI; Colavita et al. 1999), an instrument with intermediate baselines on a variety of sky projections, suitable for exploration of possible unseen nearby luminous (stellar) companions. The PTI has been demonstrated to be sensitive to nearby companions with $\Delta K < 4.0$ (Boden et al. 1998), which for a K2–3 V primary star rules out any M dwarf companions (Bessell & Brett 1988).

PTI observed HD 189733 in the K band on the nights of 2006 June 10–12, 2006 June 24, and 2006 July 8–10. Four of those nights used PTI's 85 m northwest baseline configuration, two used the 110 m north-south baseline, and one night was




FIG. 3.—SED fits of HD 189733. The top panel shows the K0 V photometric fit, the middle panel shows the K1 V fit, and the bottom panel shows the K2 V fit. All three fits allowed the reddening factor to vary.

TABLE 2
SED FITS FOR HD 189733 PHOTOMETRY TO EMPIRICAL SPECTRAL TEMPLATES

MODEL PARAMETERS ^a		FITTED PARAMETERS ^b			
Spectral Type	T_{eff} (K)	χ^2 PDF	Reddening A_V (mag)	F_{bol} (10^{-8} ergs cm^{-2} s^{-1} μm^{-1})	θ (mas)
K0 V	5188 \pm 50	2.62	0.308 \pm 0.031	3.207 \pm 0.061	0.365 \pm 0.008
K1 V ^c	5040 \pm 70	1.94	0.105 \pm 0.031	2.828 \pm 0.049	0.363 \pm 0.011
K2 V	4887 \pm 50	2.39	0.00 \pm 0.030	2.613 \pm 0.043	0.371 \pm 0.008

^a Models from Pickles (1998); T_{eff} is the effective temperature.

^b F_{bol} is the bolometric flux, and θ is the angular diameter.

^c K1 V model interpolated from the K0 V and K2 V models.

a 85 m southwest baseline. For all of these nights, HD 189733's normalized V data points were indistinguishable from unit visibility, which corresponds to a completely unresolved point source, as would be expected for a single ~ 0.37 mas star being observed by PTI at $2.2 \mu\text{m}$.

5. CONCLUSION

Our results for the radii of the host star and planet in the HD 189733 exoplanet system are formally in good agreement with existing measurements of these parameters as well as with the estimate for the density of the planet, and they have the additional and significant merit that they represent *direct* measurements of stellar and planetary diameters that do not rely on inferences about stellar atmospheres. While the diameter measurements are currently at a 6% level of accuracy, we expect to improve this considerably as we implement fringe detection at shorter wavelengths at the CHARA Array. In the meantime, these results demonstrate a new role that long-baseline optical/infrared interferometry can play in the field of exoplanet astronomy.

We would like to thank Andy Boden for sharing his SED fit tools with us, which we used to produce the fits seen in

Figures 1 and 3, and we appreciate the care that CHARA Array operator P. J. Goldfinger used in obtaining many of these observations. The CHARA Array is funded by the National Science Foundation through NSF grants AST-0307562 and AST-06006958 and by Georgia State University through the College of Arts and Sciences and the Office of the Vice President for Research. Observations with PTI are made possible through the efforts of the PTI Collaboration, which we acknowledge. Funding for PTI was provided to the Jet Propulsion Laboratory under its TOPS (Towards Other Planetary Systems), ASEPS (Astronomical Studies of Extrasolar Planetary Systems), and Origins programs and from the JPL Director's Discretionary Fund. Part of the work described in this Letter was performed at the Jet Propulsion Laboratory under contract with the National Aeronautics and Space Administration. This research has made use of the SIMBAD literature database, operated at CDS, Strasbourg, France, and of NASA's Astrophysics Data System. This publication makes use of data products from the Two Micron All Sky Survey (2MASS), which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

REFERENCES

- Bakos, G. Á., et al. 2006a, *ApJ*, 641, L57
 ———. 2006b, *ApJ*, 650, 1160
 Baraffe, I., et al. 1998, *A&A*, 337, 403
 Bessell, M. S. 2000, *PASP*, 112, 961
 Bessell, M. S., & Brett, J. M. 1988, *PASP*, 100, 1134
 Boden, A. F., et al. 1998, *ApJ*, 504, L39
 Bouchy, F., et al. 2005, *A&A*, 444, L15
 Charbonneau, D., et al. 2000, *ApJ*, 529, L45
 Claret, A., Diaz-Cordoves, J., & Gimenez, A. 1995, *A&AS*, 114, 247
 Colavita, M. M., et al. 1999, *ApJ*, 510, 505
 Cutri, R. M., et al. 2003, The IRSA 2MASS All-Sky Point Source Catalog (Pasadena: NASA/IPAC), <http://irsa.ipac.caltech.edu/applications/Gator/>
 Girardi, L., et al. 2002, *A&A*, 391, 195
 Hanbury Brown, R., et al. 1974, *MNRAS*, 167, 475
 Kervella, P., et al. 2004, *A&A*, 426, 297
 Masana, E., Jordi, C., & Ribas, I. 2006, *A&A*, 450, 735
 McAlister, H. A., et al. 2005, *ApJ*, 628, 439
 Munari, U., et al. 2005, *A&A*, 442, 1127
 Olsen, E. H. 1993, *A&AS*, 102, 89
 Perryman, M. A. C., et al. 1997, *A&A*, 323, L49
 Pickles, A. J. 1998, *PASP*, 110, 863
 Sato, B., et al. 2005, *ApJ*, 633, 465
 ten Brummelaar, T. A., et al. 2005, *ApJ*, 628, 453
 van Belle, G. T., & van Belle, G. 2005, *PASP*, 117, 1263
 Winn, J. N., et al. 2007, *AJ*, 133, 1828