

Observations of Trans-Neptunian Objects at True Opposition

A. Verbiscer (1), N. Chanover (2), J. Holtzman (2), A. Hagen (3), L. Zernow (3), K. Orr (4)


(1) University of Virginia, Charlottesville, VA (verbiscer@virginia.edu) (2) New Mexico State University, Las Cruces, NM (3) Northern Arizona University, Flagstaff, AZ (4) Baylor University, Waco, TX

With observations at true opposition and at larger solar phase angles, we measure the opposition surge of several TNOs in multiple bandpasses including R , I , and occasionally V . We survey binary TNOs as well as those not currently known to have multiple components, from a variety of dynamical classes. We test the hypothesis that ejecta exchange [1] enhances the opposition surge of binary TNOs [2] by creating a surface with a complex microtexture conducive to the constructive interference which produces the effect at the smallest phase angles.

Most airless, particulate surfaces exhibit a dramatic, non-linear increase in reflectance as the solar phase angle decreases to zero. This is the opposition effect, or surge, a consequence of both interparticle shadow hiding and a constructive interference phenomenon known as coherent backscatter [3-6].

When observed at opposition, the large heliocentric distances of trans-neptunian objects (TNOs) present unique opportunities to probe regions of the phase function inaccessible to objects closer to the Sun. Since the Sun is not a point source, even the most distant TNOs cannot be observed at solar phase angle $\alpha = 0^\circ$. The angular size of the solar radius seen from the TNO establishes the minimum phase angle at which the object can be observed from Earth. Seen from the TNO, the Earth transits the solar disk, and in this configuration, the TNO is at “true” opposition. This precise alignment enables the measurement of its reflectance at the smallest solar phase angles for any body in the Solar System, down to $\alpha \simeq 0.003^\circ$ for the outermost objects. When combined with additional photometric observations at larger phase angles, we can measure the opposition surge slope (magnitudes/degree) which is a potential classifier of dynamical class and as such can relate the evolution of a TNO to its physical surface properties.

Our survey includes several classical, resonant, detached, and scattered objects [7] as well as Centaurs. We measure the opposition surge at multiple bandpasses to investigate any wavelength dependence of surge characteristics such as amplitude and angular width. Two binary TNOs, 1997 CS₂₉ and 2000 YW₁₃₄ exhibit remarkably strong opposition surges, with slopes of 2 magnitudes/degree out to $\alpha = 1^\circ$. The other, 1995 TL₈, shows a strong spike (1 magnitude/degree) between true opposition (the minimum phase angle) and



$\alpha = 0.1^\circ$, but then flattens considerably to only 0.1 magnitude/degree between phase angles 0.1° and 0.7° . Although each of these objects comes from a different dynamical class, 1997 CS₂₉ and 2000 YW₁₃₄ have secondaries with comparable semi-major axes, 2300 and 1900 km, respectively, while 1995 TL₈ is a much closer binary with only 420 km separating the primary and secondary.

If ejecta exchange plays a role in enhancing the opposition surge of a TNO binary, the effect is larger for separations of ~ 2000 km than it is for the more tightly bound components; however, the confirmation of any relationship between opposition surge strength, binarity, and hence semi-major axis, requires more observations.

- [1] Stern, A., 2009 *Icarus*, 199, 571-573.
- [2] Rabinowitz, D. L., et al. 2009. *BAAS*, 41, 65.09.
- [3] Hapke, B. 1990 *Icarus*, 88, 407-417.
- [4] Mishchenko, M. 1992 *J. Opt. Soc. Am.* 9, 978-982.
- [5] Muinonen, K., 1990 *Light scattering by inhomogeneous media: Backward enhancement and reversal of linear polarization*. Helsinki Univ.
- [6] Gladman, B. 2008 In *The Solar System Beyond Neptune*, Barucci et al., Eds., Univ. of Ariz. Press, Tucson, 43-57.
- [7] Rabinowitz, D. L., et al. 2009. *BAAS*, 41, 65.09.